

TG-EQUIVARIANCE OF CONNECTIONS AND GAUGE TRANSFORMATIONS

H. M. Bosselmann
N.A. Papadopoulos
Institut für Physik
Johannes Gutenberg-Universität
D-55099 Mainz, Germany

December 6, 1994

Abstract

We consider the notion of a connection on a principal bundle $P(M, G)$ from the point of view of the action of the tangential group TG . This leads to a new definition of a connection and allows to interpret gauge transformations as effect of the TG action.

1 Introduction

The central role the gauge potential plays in the theories of fundamental interactions of nature always justifies new investigations concerning its nature. If we take into account gauge transformations it becomes quite apparent that gauge potential is something special compared to any other particle field. It is not a field which transforms itself like a tensor field. It is something more complicated than a tensor field with more involved and deeper lying properties.

A gauge potential is carrier of a geometric structure in the sense of the metric structure in the general relativity, but in a more general sense. Because of its complexities, its deepness and its importance, there exist several formulations of it [1, 2]. In this paper, we would like to present a new property of the gauge potential: the TG -equivariance of a connection on a principal bundle $P(M, G)$ with M the space-time (basis manifold) and G the structure group. The G -equivariance property of a connection is of course well known. Here we mean, as we shall explain, the equivariance respectively to the bigger group, the tangential group TG , of the structure group G . It is also important to note that the G -equivariance property is shared by the usual tensor fields, considered as vector-valued functions on a principal bundle $P(M, G)$. But only the connection has in addition the TG -equivariance property.

The motivation for this paper came from considerations within the G -theory framework and applications of it in investigating the structure of the space of reducible connections for Yang-Mills theories [3], anomalies [4, 5] and recently within a non commutative geometrical approach to the standard model [6]. The G -theory point of view emphasizes particularly the action of a group and all what follows from it on a relevant space of physical objects.

Our starting point is as already mentioned a principal bundle $P(M, G)$ with a connection ω on it and the structure group G acting by definition freely on P from the right. Going now to the tangential space TP and TG of P and G respectively, it is important to realize that TG may be considered as a Lie group and that it acts freely on TP . The multiplication law $TG \times TG \rightarrow TG$ and the action $TP \times TG \rightarrow TP$, are to be defined below. This leads to the fact that the tangential space TP is also a principal bundle with basis manifold TM and structure group TG . We now consider the space TP/G as a fiber bundle (non-vector bundle) with typical fibre the Lie algebra $\mathcal{G}$ of G . We show that TP/G is isomorphic to the associated fiber bundle of $TP : TP \times_{TG} \mathcal{G}$ where the action of TG on $\mathcal{G}$ is deduced from the multiplication law in TG . We give a new definition of a connection on P : a connection on $P(M, G)$ is a linear section $\tilde{H}$ on the fiber bundle $TP \times_{TG} \mathcal{G}$

$$\tilde{H} : TM \rightarrow TP \times_{TG} \mathcal{G} . \quad (1)$$

It can be shown that ω (from the usual definition of a connection) and $\tilde{H}$, as above, are equivalent. The section $\tilde{H}$ may also be considered as a TG -equivariant and linear function $\bar{H}$ on TP with values on $\mathcal{G}$.

$$\begin{aligned} \bar{H} : TP &\rightarrow \mathcal{G} , \\ X_p &\mapsto \bar{H}(X_p) \quad \text{with} \quad \bar{H}(X_p \cdot \xi_g) = \xi_g^{-1} \cdot \bar{H}(X_p) , \end{aligned} \quad (2)$$

and $X_p \in TP$, $\xi_g \in TG$. $X_p \cdot \xi_g$ represents the action of TG on TP from the right and $\xi_g \cdot \eta$ with $\eta \in \mathcal{G}$ the action of TG on $\mathcal{G}$ (from the left). It follows immediately that this property of TG -equivariance of $\bar{H}$ is also adapted by the ω . As we shall see, the following equation is valid:

$$\omega(X_p) = -\bar{H}(X_p) . \quad (3)$$

The last topic we would like to discuss, are the gauge transformations on ω in the light of the above development. It is well known that a gauge transformation Ψ can be described by the equivariant function U on P with values on G . So we have $\Psi(p) = pU(p)$ with

$$U : P \rightarrow G$$

and

$$U(pg) = g^{-1}U(p)g . \quad (4)$$

This makes the G -action point of view transparent and induces the gauge transformation of the connection ω given by:

$$(\Psi^*\omega)(X_p) = \text{Ad}_{U(p)^{-1}}\omega(X_p) + U^{-1}dU(X_p) . \quad (5)$$

This transformation may also be expressed by the equation

$$(\Psi^*\omega)(X_p) = -(TUX_p)^{-1} \cdot (-\omega(X_p)) . \quad (6)$$

As we see, gauge transformations expressed pointwise are nothing else but the TG -action on the values of connection. The non-linear term in the transformation of a connection may now be explained by the TG -action on $\mathcal{G}$. TG is a group of affine type which acts non linearly on $\mathcal{G}$.

In what follows we discuss in the next section the multiplication rule in TG , the action of TG on TP and the space TP as a principal bundle over TM . In section three we present the new definition of a connection and focus on its TG -equivariance property. In section four we consider gauge transformations on a connection from the point of view of the TG -action and we finally give a summary in section five.

In the physical literature we found no explicit treatment of the TG -equivariance aspect of connections. The same is valid, to our surprise, for the recent mathematical literature we examined. The only place we found some, for us very inspiring, but short and compact indications on TG -equivariance, is the very early paper [7]. Parts of section two and three are implicitly or explicitly contained in [7] but without connection to the gauge transformations which were the main motivation for the present paper. For all these reasons we tried to be quite detailed and explicit in our presentations.

2 TP as TG principal bundle

The action of the Lie-group G in the principal bundle $P(M, G)$ is a part of its definition. Therefore we expect the G -action to be also relevant for further structures defined on P . This is the case i.e. for the connection ω on P and for the vector valued functions on P which correspond to particle fields in the space time M .

In this paper we would like to investigate the action of the group TG which of course contains G and in addition its Lie algebra $\mathcal{G} = \text{Lie}(G)$. For this purpose we have first to determine the multiplication law on TG and then the action of TG on TP . For the reasons explained in the introduction, we shall proceed in a quite detailed way. We start with the multiplication law in TG , although the space TG is isomorphic to the Cartesian product of $\mathcal{G}$ and G , the relevant multiplication law is not the so expected trivial one, but it makes TG a semi-direct product of $\mathcal{G}$ and G :

$$\begin{aligned} TG \times TG &\rightarrow TG, \\ (\xi_g, \eta_h) &\mapsto \xi_g \cdot \eta_h := TR_h\xi_g + TL_g\eta_h . \end{aligned} \quad (7)$$

The notation R and L is used to the right and left action of G on G . The inverse of ξ_g is given by

$$\xi_g^{-1} = -TL_{g^{-1}}TR_{g^{-1}}\xi_g . \quad (8)$$

If we denote with e and 0_e the neutral elements in G and TG we obtain by explicit calculation $\xi_g \cdot (\xi_g)^{-1} = 0_e$. We have of course the injections:

$$\begin{array}{ccc} G & \rightarrow & TG \\ g & \mapsto & 0_g \end{array} \quad \text{and} \quad \begin{array}{ccc} \mathcal{G} & \rightarrow & TG \\ \xi & \mapsto & \xi_e \end{array} . \quad (9)$$

The verification of associativity is slightly involved and it is therefore an advantage to use the isomorphism which contains the semi-direct product splitting of TG given by

$$\begin{aligned} \alpha : TG &\rightarrow \mathcal{G} \otimes G , \\ \xi_g &\mapsto (\xi, 0_g) := (TR_{g^{-1}}\xi_g, g) . \end{aligned} \quad (10)$$

Denoting as usual $\text{Ad}_G = \text{Aut}(\mathcal{G})$ with

$$\text{Ad}_g \xi = TL_g TR_{g^{-1}} \xi = 0_g \xi 0_{g^{-1}} , \quad (11)$$

we have for the multiplication rule the multiplication

$$\xi_g \cdot \eta_h \mapsto (\xi, g)(\eta, h) = (\xi + \text{Ad}_g \eta, gh) . \quad (12)$$

At this point it should be noticed that of course all the above structures are induced naturally by the T -functor and the canonical isomorphism given in eq. (10)

In what follows we need several times the action of TG on $\mathcal{G}$. It may be derived from the multiplication law above and the action of TG on the G -orbit space TG/G given by

$$TG/G = \{\xi_g G = TR_h \xi_g | h \in G\} \quad (13)$$

which we also denote as $\mathcal{G} = TG/G$ by misuse of notation. The rule of addition in TG/G may be defined by

$$\xi_g G + \eta_g G := (\xi_g + \eta_g)G . \quad (14)$$

The TG -action on $\mathcal{G} = TG/G$ is given by¹

$$\begin{aligned} TG \times \mathcal{G} &\rightarrow \mathcal{G} , \\ (\eta_h, \xi) &\mapsto \eta_h \cdot \xi = \text{Ad}_h \xi + TR_{h^{-1}} \eta_h . \end{aligned} \quad (15)$$

¹The restriction to the subgroup G of TG leads as expected to the adjoint action of G on $\mathcal{G}$:

$$\begin{aligned} G \times \mathcal{G} &\rightarrow \mathcal{G} \\ (g, \eta) &\mapsto TR_{g^{-1}}(0_g \cdot \eta) = TR_{g^{-1}} TL_g \eta = \text{Ad}_g \eta \end{aligned}$$

The action of TG on TP is given canonically from the lifting of the G -action on P to the tangential space (T -functor). So we have in an obvious notation from

$$\begin{aligned} R : P \times G &\rightarrow P, \\ (p, g) &\mapsto pg = R_g p = R_p g, \end{aligned} \quad (16)$$

the TG -action on TP :

$$\begin{aligned} TP \times TG &\rightarrow TP, \\ (X_p, \xi_g) &\mapsto X_p \cdot \xi_g := TR_g X_p + TR_p \xi_g. \end{aligned} \quad (17)$$

In order to facilitate the reading, we give some further notations and some examples. For the Maurer–Cartan form C on G given by

$$C(\xi_g) = TL_{g^{-1}} \xi_g. \quad (18)$$

we often use the expression $\xi(g) = C(\xi_g) \in \mathcal{G}$. We denote a fundamental vector field on P with $\hat{\xi}$ where we have $\xi \in \mathcal{G}$ and $\hat{\xi}_p = TR_p \xi$.

We now give some useful examples of the TG action on TP .

$$\begin{aligned} X_p \cdot 0_e &= X_p, \\ X_p \cdot \xi &= X_p + \hat{\xi}_p, \\ X_p \cdot 0_g &= TR_g X_p, \\ 0_p \cdot \xi &= TR_p \xi =: \hat{\xi}_p, \\ 0_p \cdot \xi_g &= TR_{pg} C(\xi_g) =: TR_{pg} \xi(g). \end{aligned} \quad (19)$$

After this preparation, everything is fixed and as a first result of the TG -action we may formulate the following proposition:

The space TP is a principal bundle $TP(TM, TG)$ which structure group TG basis manifold TM and canonical projection $T\pi : TP \rightarrow TM$. π is the canonical projection in $P(M, G)$, $\pi : P \rightarrow M$.

This can be seen from the fact that

- i) TG acts freely on TP .
- ii) $T\pi$ commutes with the group action TG .

For i) we have to show that with $X_p \cdot \xi_g = X_p$,

$$\xi_g = 0_e \quad (20)$$

must be valid.

From

$$X_p \cdot \xi_g = TR_g X_p + TR_p \xi_g = X_p \quad (21)$$

we must first have $g = e$ so that using eq. (19), we obtain

$$X_p \cdot \xi_e = X_p + \hat{\xi}_p = X_p \quad \text{and} \quad \hat{\xi}_p = 0_p \quad (22)$$

which shows that

$$\xi = 0 \quad \text{and} \quad \xi_g = 0_e . \quad (23)$$

For ii) we have to show that

$$T\pi X_p = T\pi(X_p \cdot \xi_g) \quad (24)$$

is valid. Using eq. (19) and

$$X_p \cdot \xi_g = X_p \cdot 0_g + 0_p \cdot \xi_g \quad (25)$$

we obtain

$$X_p \cdot \xi_g = TR_g X_p + TR_{pg} \xi(g) . \quad (26)$$

$TR_{pg} \xi(g)$ is vertical and

$$T\pi TR_g X_p = T\pi X_p , \quad (27)$$

so that ii) is indeed valid.

3 The TG -equivariance of connection

Having in mind the G -equivariance property of connections and the TG -action on TP , as discussed in the previous section, we study the space TP/G and ask ourselves what its connection with the connections on $P(M, G)$ is. This leads us to a new definition of a connection which is of course, as we shall show, equivalent to the usual one. The original definition of connections is closely related to the G -action aspect whereas the definition in this section will reveal the TG -action aspect and bring to light the most characteristic property of connections: the TG -equivariance.

For that purpose we shall first discuss the G orbit space

$$TP/G = \{X_p G = \{TR_g X_p | g \in G\} | p \in P\} \quad (28)$$

and we shall show that it is isomorph to the TP associated fibre bundle with typical fiber the space $\mathcal{G}$:

$$TP/G \simeq TP \times_{TG} \mathcal{G} . \quad (29)$$

The TG -action on $\mathcal{G}$ is given in eq. (15). This action is non-linear, therefore we do not expect $TP \times_{TG} \mathcal{G}$ to be a vector bundle although it may be represented as $TP \times_{TG} \mathcal{G} = TM \tilde{\times} \mathcal{G}$.

The isomorphism in eq. (29) follows from the fact that TP is a principal bundle and that G is a subgroup of its structure group TG :

$$\begin{aligned} TP/G &\rightarrow TP \times_{TG} (TG/G) , \\ [X_p \cdot \xi_g]_G &\mapsto [X_p, \xi_g G]_{TG} . \end{aligned} \quad (30)$$

If we take $\xi_g = 0_e$, this may be seen more perspeciously:

$$X_p G \mapsto [X_p, 0_e G]_{TG} . \quad (31)$$

In the fibre bundle TP/G , we have to define an addition: for $\pi(p) = \pi(q)$ and $p = qg$, we have:

$$X_p G + Y_q G := X_p G + (TR_g Y_q)G = (X_p + TR_g Y_q)G . \quad (32)$$

In what follows we shall give three new definitions of a connection. These are closely related to the horizontal lifting of the (original) definition of connection: *A connection on a principal bundle $P(M, G)$ is a linear section in the associated fibre bundle $TP \times_{TG} \mathcal{G}$ ($\mathcal{G} = \text{Lie}(G)$).*

The action of TG on $\mathcal{G}$ is given in eq. (15). If we denote by $\tilde{H}$ this section, we have:

$$\begin{aligned} \tilde{H} : TM &\rightarrow TP/G , \\ V_x &\mapsto \tilde{H}(V_x) . \end{aligned} \quad (33)$$

The existence of such a section leads to an equivalent but more abstract definition:

A connection on a principal bundle $P(M, G)$ is a reduction of the structure group TG of the principal bundle $TP(TM, TG)$ to G .

The most transparent definition we obtain if we consider the above section $\tilde{H}$ in TP/G as a $\mathcal{G}$ valued linear function $\bar{H}$ on TP :

A connection on a principal bundle $P(M, G)$ is a linear TG -equivariant function on TP with values in $\mathcal{G} = \text{Lie}(G)$. $\bar{H}$ is given by

$$\begin{aligned} \bar{H} : TP &\rightarrow \mathcal{G} , \\ X_p &\mapsto \bar{H}(X_p) \quad \text{with} \\ \bar{H}(X_p \cdot \xi_g) &= (\xi_g)^{-1} \cdot \bar{H}(X_p) . \end{aligned} \quad (34)$$

At this point it is interesting to note that the function $\bar{H}$ is, as we shall show, directly related to the original definition of a connection ω (as a linear function on TP) whereas the section $\tilde{H}$ is directly related to the horizontal lift H^ω which stems from the ω . The relation between $\tilde{H}$ and $\bar{H}$ is given by:

$$\tilde{H}(T\pi X_p) = [X_p, \bar{H}(X_p)]_{TG} . \quad (35)$$

The linearity of $\tilde{H}$ is related as expected to the linearity of $\tilde{H}$:

$$\begin{aligned}
[X_p, \xi]_{TG} + [Y_p, \eta]_{TG} &= [X_p + \hat{\xi}_p, 0]_{TG} + [Y_p + \hat{\eta}_p, 0]_{TG} , \\
&= [X_p \cdot \xi, 0]_{TG} + [Y_p \cdot \eta, 0]_{TG} , \\
&= (X_p + \hat{\xi}_p)G + (Y_p + \hat{\eta}_p)G , \\
&= (X_p + \hat{\xi}_p + Y_p + \hat{\eta}_p)G , \\
&= [X_p + Y_p + \hat{\xi}_p + \hat{\eta}_p, 0]_{TG} , \\
&= [X_p + Y_p, \xi + \eta]_{TG}
\end{aligned} \tag{36}$$

where we have used the isomorphism of eq. (31) and the definition of the eq. (32).

This corresponds to

$$[X_p, \bar{H}(X_p)]_{TG} + [Y_p, \bar{H}(Y_p)]_{TG} = [X_p + Y_p, \bar{H}(X_p) + \bar{H}(Y_p)]_{TG} . \tag{37}$$

So we obtain from the linearity of $\tilde{H}$

$$\tilde{H}(T\pi X_p) + \tilde{H}(T\pi Y_p) = \tilde{H}(T\pi(X_p + Y_p)) \tag{38}$$

the linearity of $\bar{H}$:

$$\begin{aligned}
[X_p, \bar{H}(X_p)]_{TG} + [Y_p, \bar{H}(Y_p)]_{TG} &= [X_p + Y_p, \bar{H}(X_p + Y_p)]_{TG} \\
\bar{H}(X_p + Y_p) &= \bar{H}(X_p) + \bar{H}(Y_p) .
\end{aligned} \tag{39}$$

For the relation between the horizontal lift H^ω and $\tilde{H}$, taking $\pi(p) = x$ and $T\pi X_p = V_x$, we have:

$$\begin{aligned}
H^\omega_p : T_x M &\rightarrow T_p P , \\
V_x &\mapsto H^\omega_p(V_x) .
\end{aligned} \tag{40}$$

Using the projection:

$$\begin{aligned}
\mu : TP &\rightarrow TP/G , \\
X_p &\mapsto X_p G ,
\end{aligned} \tag{41}$$

we have

$$\tilde{H}_x = \mu \circ H^\omega_p . \tag{42}$$

The relation of $\bar{H}$ to connection ω is given by

$$\omega(X_p) = -\bar{H}(X_p) . \tag{43}$$

All we have to show is that the so-defined $\omega = \omega(\bar{H})$ fulfils the original axioms of a connection:

- i) ω is G -equivariant.
- ii) the restriction of ω to the fundamental vector fields on P corresponds, after the identification of the fundamental vector fields with the Lie algebra $\mathcal{G}$, to the identity.

The condition i) is clearly valid since the ω given by $\bar{H}$ is TG -equivariant, G is a subgroup of TG and acts on $\mathcal{G}$ like Ad_G .

The condition ii) follows from the fact that for $\hat{\xi}_p = TR_p\xi$ with $\xi \in \mathcal{G}$

$$\begin{aligned}
\omega(\hat{\xi}_p) &= -\bar{H}(\hat{\xi}_p) = -\bar{H}(0_p \cdot \xi_e) , \\
&= -\xi_e^{-1} \cdot \bar{H}(0_p) , \\
&= -\xi^{-1} \cdot 0 , \\
&= -(-\xi) = \xi
\end{aligned} \tag{44}$$

where we have used the TG -equivariance of $\bar{H}$ (eq. (34)) and the linearity of $\bar{H}$ (eq. (39)) to evaluate $\bar{H}(0_p)$.

The above considerations showed the equivalence of all the notions of connection represented by ω , H^ω , $\tilde{H}$, $\bar{H}$ in every combination and particularly in (ω, H^ω) with $(\bar{H}, \tilde{H})$.

At this point we give, for those who feel uneasy with the use of the fibre bundle $(TP/G \rightarrow TM) = TP \times_{TG} \mathcal{G}$ in the definition of a connection in eq. (33), a further definition in terms of vector bundles. For that purpose we observe that $TP/G \rightarrow M$ is a vector bundle with typical fiber $\mathbf{R}^m \times \mathcal{G}$ and $m = \dim(M)$. So, denoting this vector bundle by $E = (TP/G \rightarrow M)$, we have $E = M \tilde{\times} (\mathbf{R}^m \times \mathcal{G})$.

This leads to the definition of a connection as a special vector bundle homomorphism between TM and E :

A connection on a principal bundle $P(M, G)$ is a vector bundle homomorphism $\hat{H}$:

$$\hat{H} : TM \rightarrow E = (TP/G \rightarrow M) \tag{45}$$

with the restriction $\nu \circ \hat{H} = \text{id}_{TM}$ and $\nu = T\pi \circ \mu^{-1}$ the projection $\nu : TP/G \rightarrow TM$. In the case of a trivial $P(M, G)$, this definition corresponds to the graph of a gauge potential A :

$$A : TM \rightarrow \mathcal{G} \tag{46}$$

as it is used in particle physics phenomenology.

4 TG -equivariance and gauge transformations

In this section we deal with the implications for gauge transformations from the TG -action point of view. In this way we obtain a new formulation for

the well known expression for gauge transformations of connections in terms of TG -elements instead of G -elements.

We consider a gauge transformation as an automorphism on $P(M, G)$ which may be expressed equivalently by a function on P with values on G [8]:

$$\begin{aligned}\Psi : P &\rightarrow P, \\ p &\mapsto \Psi(p) = pU(p)\end{aligned}\tag{47}$$

and

$$\begin{aligned}U : P &\rightarrow G, \\ p &\mapsto U(p)\end{aligned}$$

with

$$U(pg) = g^{-1}U(p)g.\tag{48}$$

The automorphism Ψ induces in the space of connections the transformation Ψ^* and we have for the connection ω pointwise the expression

$$(\Psi^*\omega)(X_p) = \text{Ad}_{U(p)^{-1}}\omega(X_p) + (U^*C)(X_p)\tag{49}$$

with $\xi_p \in TP$ and where C is the Maurer–Cartan form on G :

$$\begin{aligned}C : TG &\rightarrow \mathcal{G}, \\ \xi_g &\mapsto C(\xi_g) = TL_{g^{-1}}\xi_g.\end{aligned}\tag{50}$$

We would like now to reexpress the right hand side of eq. (49) in terms of the TG -action on TP and $\mathcal{G}$. For the action of TG on $\mathcal{G}$ we obtain from

$$\xi_g \cdot \eta = \text{Ad}_g\eta + TR_{g^{-1}}\xi_g,$$

given in eq. (15), the $(\xi_g)^{-1}$ action on η :

$$(\xi_g)^{-1} \cdot \eta = \text{Ad}_{g^{-1}}\eta - TL_{g^{-1}}\xi_g\tag{51}$$

or

$$(\xi_g)^{-1} \cdot \eta = \text{Ad}_{g^{-1}}\eta - C(\xi_g).\tag{52}$$

For the $\Psi^*\omega$, expressed in $\bar{H}$, we have pointwise:

$$(\Psi^*\omega)(X_p) = \omega(T\Psi X_p) = -\bar{H}(T\Psi X_p).\tag{53}$$

Using the equivariance property of $\bar{H}$ (eq. (34)) and eq. (52), we obtain:

$$\bar{H}(T\Psi X_p) = \bar{H}(X_p \cdot TUX_p) = (TUX_p)^{-1} \cdot \bar{H}(X_p)\tag{54}$$

and

$$(TUX_p)^{-1} \cdot \bar{H}(X_p) = \text{Ad}_{U(p)^{-1}} \bar{H}(X_p) - C(TUX_p) . \quad (55)$$

Eqs. (53 - 55) lead to

$$(\Psi^* \omega)(X_p) = -(TUX_p)^{-1} \cdot (-\omega(X_p)) \quad (56)$$

or equivalently to

$$(\Psi^* \omega)(X_p) = \text{Ad}_{U(p)^{-1}} \omega(X_p) + U^* C(X_p) . \quad (57)$$

On the right hande side of eq. (56), the minus in front of $\omega(X_p)$ cannot be eliminated because the TG -action is not linear. Notice also that we obtained in this way eq. (49) in a very straightforward manner.

5 Summary

The aim of this paper was to set some new lights on several aspects of the notion of a connection on a principal bundle $P(M, G)$ and especially on its gauge transformations. We expect this to be very useful in physical applications.

We proceeded essentially by exploring the properties of the T -functor, considering the action of the tangential group TG on $\mathcal{G} = \text{Lie}(\mathcal{G})$ and TP . It was perceived that TP is also a principal bundle $TP(TM, TG)$ and that the fibre bundle $TP/G \rightarrow TM$ associated with it ($TP \times_{TG} \mathcal{G}$) is useful for a new definition of a connection as a linear section on it.

This definition revealed the most characteristic property of a connection, the TG -equivariance:

$$\omega(X_p \cdot \xi_g) = -(\xi_g)^{-1} \cdot (-\omega(X_p)) \quad (58)$$

where we have used the symbol $\cdot$ for the TG -action on TP and $\mathcal{G}$.

The TG -equivariance property of the connection also gives the most plausible explanation for the specific form the expression for gauge transformations takes. In particular, the nonlinearity in those transformations stems from the nonlinear action of the affine group TG .

Acknowledgements

We would like to thank S. Klaus and M. Kreck for very helpful discussions and careful reading of the manuscript. We also thank R. Coquereaux for an interesting discussion, and R. Buchert and F. Scheck for useful comments.

References

1. S. Kobayashi and K. Nomizu, Foundations of differential geometry, Volume 1, John Wiley & Sons: New York, London, Sydney, 1963.
2. Y. Choquet–Bruhat, C. De Witt–Morette and M. Dillard–Bleik, Analysis, Manifolds, and Physics, North–Holland: Amsterdam, Oxford, New York, Tokyo, 1987.
3. – A. Heil, A. Kersch, N. A. Papadopoulos, B. Reifenhäuser and F. Scheck, Structure of the Space of Reducible Connections for Yang–Mills Theories, Journal of Geometry and Physics: Vol. 7, n. 4, 1990, 489.
– A. Heil, A. Kersch, N.A. Papadopoulos and B. Reifenhäuser, Abelian Charges in Nonabelian Yang–Mills Theory from the Stratification of the Space of Gauge Potentials, Annals of Physics, Vol. 217, No. 2 1992.
4. – A. Heil, A. Kersch, N.A. Papadopoulos, B. Reifenhäuser, F. Scheck and H. Vogel, Anomalies from the Point of View of G–Theory, Journal of Geometry and Physics: Vol. 6, n. 2, 1989, 237.
– A. Heil, A. Kersch, N.A. Papadopoulos, E. Reifenhäuser and F. Scheck: Anomalies from Nonfree Action of the Gauge Group, Annals of Physics: 200, 1990, 206.
5. N.A. Papadopoulos, The Role of Stratification of Anomalies, in J. Debrus and A.C. Hirshfeld (Eds.), Geometry and Theoretical Physics, Springer-Verlag: Berlin, Heidelberg, 1991, 210.
6. R. Häußling, N.A. Papadopoulos and F. Scheck, Supersymmetry in the Standard Model of Electroweak Interactions, Physics Letters B303, 1993, 265.
7. S. Kobayashi, Theory of connection, Annali di Matematica, 43, 1957, 199.
8. M. Göckeler and T. Schücker, Differential Geometry, Gauge Theories, and Gravity, Cambridge University Press: Cambridge, 1987.